

ALBENKYX L



Veterinary Use



Internal Antiparasitic - Premix for oral use in feed

FORMULA:

1 g of product contains:

Albendazol	100 mg
Levamisol HCl	20 mg
Excipients q. to	1 g

DESCRIPTION:

Albenkx L is an internal antiparasitic that also acts as an immunoregulator, with ovicidal power, sustained protection and easy administration.

PRESCRIPTION:

Albenkx L is a sustained and broad-spectrum internal antiparasitic for the treatment of pigs and cattle against nematodes, flukes (fasciola hepatica), cestode infestation, and mature and immature cases of gastrointestinal nematodes (ascaris suum, strongyloides ransomi, oesophagostomun dentatum, etc.)) and pulmonary (metastrongylus sp.); Verminous gastroenteritis, parasitic bronchopneumonia. Albenkx L is the indicated treatment against infestation with different parasites in birds such as ascaris, heterakis, tapeworms, capillaries, etc.

DOSAGE AND USE:

Oral use, mixed with feed, according to the following rates:

Cattle: Gastrointestinal and pulmonary helminths 5 mg / kg / b.w. Ostertagia type II: 7.5 mg / kg / b.w. Hepatic fasciola: 10 mg / kg / b.w. Treatment duration: 5 days continuously. Administer as sole feeding.

Pigs: 15 mg / kg / body weight / day.

Poultry: 1 kg of Albenkx L in 1000 kg of feed for 7 consecutive days. Administer as sole feed. It is advisable to repeat the treatment 15 days after the first performance to avoid parasitic reinfestation.

It is advisable to mix the quantity of product to be used in an adequate quantity of feed (25 kg) before adding it to the final volume of the feed portion.

Consult the veterinary doctor to adjust the dose according to the needs of the case to be treated and according to the daily consumption of the flock.

PRECAUTIONS:

Keep out of the reach of children and pets. Avoid inhalation and direct contact of the product with the skin and eyes. If accidental exposure occurs, wash with running water and soap. Store in its original packaging in a cool, dry place between 15° and 25° C, protected from sunlight and moisture. People with known hypersensitivity to the drug should avoid contact with the product. Do not throw empty containers in open landfills, dispose of them in accordance with local regulations. Maintain adequate ventilation when administering the product. Close the container tightly after each use. Validity after being mixed with food: 15 days. Do not use after the expiration date on the label.

Contraindications: known hypersensitivity to albendazole and / or levamisole.

Side effects: occasional gastrointestinal disorders, vomiting, diarrhea, hypersalivation, foam in the mouth, pulmonary edema, dyspnea, bronchospasm, tremors, nervousness, ataxia (lack of coordination of movements), hyperesthesia, weakness, collapse, seizures. These undesirable effects generally disappear within 1 to 2 hours after administration. If you notice serious side effects or other effects not mentioned in this notice, tell your vet.

Albenkx L has no contraindications or side effects at the recommended doses.

WITHDRAWAL PERIOD: Birds: 7 days. Pigs: 5 days. Cattle: 15 days. Milk: 4 days (8 milkings).

Do not use it in the first 45 days of gestation, or within 45 days after delivery. Do not use in laying hens producing eggs for human consumption. Do not use the product in breeding hens producing fertile eggs.

PACKING:

1kg, 5kg and 20kg.

(It may be possible not all packings are traded).

