

# CEFTIOKYYX

## Antibiotic - Sterile Injectable Suspension

### FORMULA:

1 ml of product contains:  
Ceftiofur (as hydrochloride)..... 50 mg  
Vehicle q.s.to..... 1 ml



### DESCRIPTION:

CEFTIOKYYX is an antibiotic sterile suspension ready to use by parenteral means. Each ml. contains Ceftiofur Hydrochloride equivalent to 50 mg of Ceftiofur.

CEFTIOKYYX offers fast results as after its administration an improvement can be observed in the clinical state of animals as well as a limit to the pathological proliferation to other animals. It also normalizes the animal temperature since the first day of treatment. Likewise, it is observed a recovery in the consumption of feed, minimizing the loss of weight.

### PRESCRIPTION:

This product is indicated for the treatment of bacterial origin pneumonias and foot rot in bovines, bacterial pneumonia or pleuropneumonia in porcines.

### SUSCEPTIBLE ORGANISMS:

Gram negative: Pasteurella multocida, Pasteurella haemolítica, Haemophilus somnus

Bacteroides fragilis (group), Bacteroides spp, Bacteroides melaninogenicus, Actinobacillus pleuropneumoniae, Escherichia coli, Moraxella bovis, Salmonella cholerasuis, Salmonella typhimurium, Haemophilus parasuis, Actinobacillus suis.

Gram positive: Staphylococcus hyicus, Streptococcus suis, Peptostreptococcus anaerobius beta-haemolytic, Streptococcus spp

### DOSAGE AND USE:

Bovines: 1 to 2,2 ml. of suspension per each 50 kg of weight by intramuscular or subcutaneous injection. This dosage is equivalent to 1,0 - 2.2 mg of Ceftiofur per kg. of body weight. Porcines: 0,6 to 1 ml per each 10 kg. of weight by intramuscular injection, which is equivalent to 3,0 - 5,0 mg. of Ceftiofur per kg. of body weight. Treatment should be repeated every 24 hours for three days in a row. The duration of the treatment can be extended according to the Veterinary Surgeon's judgment.

### WARNINGS:

Keep out of the reach of children. Do not administer to animals with proved sensitivity to the drug. Use sterile instrumental to remove product from container. Avoid contact with the product during its use. If accidental exposure occurs, wash with tap water and soap. Do not administer by endovenous means in pigs. Only by intramuscular means. Wash hands and change clothes once product has been administered. Store at temperatures between 15° and 25° C. Do not freeze. Shake well before use.

### Use restrictions:

Withdrawal period in Bovines: 48 hours. In pigs: 5 days.

### PACKING:

50 ml and 100 ml glass vials.

