

CIPROKYYX 200



Chemoterapheutic - Antimicrobial - Solution for oral use

Veterinary use



FORMULA:

Each ml contains:
Ciprofloxaxine..... 200 mg
Excipients q. to..... 1 ml

DESCRIPTION:

CIPROKYYX 200 is an antibiotic solution based on ciprofloxacin which belongs to the group of quinolones and works with bactericide action against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp.

PRESCRIPTION:

CIPROKYYX 200 is indicated for the prevention and treatment of gastrointestinal, respiratory and urinary tract infections caused by ciprofloxacin sensitive micro-organisms like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp. in calves, goats, poultry, sheep and swine.

DOSAGE AND USE:

For oral administration diluted in drinking water.

Calves, goats and sheep: Twice daily 5 ml every 75 - 150 kg of body weight for 3 to 5 days.

Poultry: At the rate of 10 mg/kg b.w. Approximately 100 ml every 300 - 400 litres of drinking water for 3 to 5 days.

Swine: At the rate of 10 mg/kg b.w. Approximately 100 ml every 200 - 600 litres of drinking water for 3 to 5 days.

Note: for pre-ruminant calves, lambs and kids only.

Only sufficient medicated drinking water should be prepared to cover daily requirements. Medicated drinking water should be replaced every 24 hours.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS:

Keep out of the reach of children. Avoid direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Use suitable and protective garment when administering medication. Wash hands and change clothes after administration. Store in its original container at temperatures between 15°C and 25°C avoiding direct sunlight on stored product. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Do not freeze. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Do not use after expiry date which is stated on the label.

Contraindications: Hypersensitivity to ciprofloxacin. Administration to animals with a serious impaired hepatic and/or renal function. Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

Side effects: Hypersensitivity reactions. Administration to juvenile animals can lead to arthropathy.

CYPROKYYX 200 has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawal period for meat: 12 days.

Do not use in laying hens producing eggs for human consumption.

PACKING:

500 ml and 1 litre HDPE bottle.

May be possible that not all packages are traded.

