

FLOXON 100

Chemotherapeutic - Antimicrobial - Solution for oral use



FORMULA:

100 ml of product contain:

Enrofloxacin..... 10 g
Vehicle q.s.to..... 100 ml

DESCRIPTION:

FLOXON 100 is a solution that consists of ENROFLOXACIN, a broad spectrum chemotherapeutic of the quinolone carboxyl acids family, a synthesis product developed especially for Veterinary Medicine. Due to its bactericidal effect on Gram positive, Gram negative bacteria and mycoplasma, produces a fast decline of avian mortality. It also produces a fast decrease in the symptoms preventing relapses.

PRESCRIPTION:

FLOXON 100 is especially indicated for the treatment of avian infectious diseases produced by Gram positive bacteria, Gram negative bacteria and mycoplasma (*M. gallisepticum*; *M. synoviae*; *M. meleagridis*). Indicated for the oral treatment of broilers, replacement pullets and breeders.

ENROFLOXACIN ANTIBACTERIAL ACTIVITY: Negative Coagulase *Staphylococcus*, *Staphylococcus aureus*, *Streptococcus pyogenes*, *Streptococcus faecalis*, *Streptococcus pneumoniae*, *Escherichia coli*, *Klebsiella* spp, *Serratia marcescens*, *Proteus mirabilis*, *Proteus vulgaris*, *Proteus stuartii*, *Providencia* spp, *Salmonella* spp, *Shigella* spp, *Pseudomonas aeruginosa*, *Haemophilus* spp, *Legionella*, *Campylobacter* spp, *Bordetella* spp, *Pasteurella* spp, *Moraxella* spp, *Clostridium difficile*, *Bacteroides* spp, *Mycoplasma*, *Ureaplasma*, *Chlamidia*.

DOSAGE AND USE:

For oral administration in drinking water, considering it being the sole supply during the medication.

Therapeutic dose is 10 mg/kg of b.w. Approximately 50 ml of product every 100 L of drinking water which is 50 p.p.p.m. Treatment duration: 3 to 5 days.

For Salmonellosis. 500 ml every 1000 L of drinking water during 5 days.

Only sufficient medicated drinking water should be prepared to cover daily requirements. Medicated drinking water should be replaced every 24 hours.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS:

Keep out of the reach of children. Avoid direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Use suitable and protective garment when administering medication. Wash hands and change clothes after administration. Store in its original container at temperatures between 15°C and 25°C avoiding direct sunlight on stored product. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Do not freeze. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Do not use after expiry date which is stated on the label.

Contraindications: Hypersensitivity to enrofloxacin. Administration to animals with a serious impaired hepatic and/or renal function. In cases of disorders in the central nervous system and in very young animals due to joint problems.

Side effects: Hypersensitivity reactions, gastrointestinal disorders, skin pruritus and / or photosensitivity, joint pain, seizures

FLOXON 100 has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawal period for poultry: 10 days. Do not use in laying hens producing eggs for human consumption.

PACKING:

100 ml and 1 Liter HDPE bottle.

May be possible that not all packages are traded.

