

# FLOXON 200B



Veterinary use



## Antibiotic - Solution for oral use

### FORMULA:

Each ml of product contains:

Enrofloxacin..... 200 mg  
Bromhexine hcl..... 15 mg  
Vehicle q.s.to..... 1 ml

### DESCRIPTION:

FLOXON 200B is a chemotherapeutic at 20 % enrofloxacin concentration combined with bromhexine, wich has a mucolytic action. It is a solution for the oral treatment of broiler chickens, breeders, 16 weeks old replacement pullets.

### PRESCRIPTION:

FLOXON 200B is specially indicated for the treatment of poultry infectious diseases in poultry, produced by Gram poistive and Gram negative bacteria and/or mycoplasmas.

**ENROFLOXACIN ANTIBACTERIAL ACTIVITY:** Negative Coagulase Staphylococcus, Staphylococcus aureus, Streptococcus pyogenes, Streptococcus faecalis, Streptococcus pneumoniae, Escherichia coli, Klebsiella spp, Serratia marcescens, Proteus mirabilis, Proteus vulgaris, Proteus stuartii, Providencia spp, Salmonella spp, Shigella spp, Pseudomonas aeruginosa, Haemophilus spp, Legionella, Campylobacter spp, Bordetella spp, Pasteurella spp, Moraxella spp, Clostridium difficile, Bacteroides spp, Mycoplasma, Ureaplasma, Chlamidia.

### DOSAGE AND USE:

For oral administration in drinking water.

**Poultry:** 25 ml of product in 100 litres of drinking water (10 mg/kg body weight) during 3 to 5 days. Salmonellosis: 5 days. Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

### WARNINGS:

Keep out of the reach of children and domestic animals. Avoid direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Use suitable and protective garment when administering medication. Wash hands and change clothes after administration. Store in its original container at temperatures between 15°C and 25°C avoiding direct sunlight on stored product. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Do not freeze. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Do not use after expiry date which is stated on the label.

**Contraindications:** anticoagulants, aluminum, cimetidine, xanthines in general. In cases of disorders in the central nervous system and in very young animals due to joint problems.

**Side effects:** gastrointestinal disorders, skin pruritus and/or photosensitivity, joint pain, seizures. If you notice any serious side effects or other effects not listed on this notice, please inform your veterinarian.

FLOXON 200B has no contraindications or side effects at recommended doses.

**Use restrictions:** Withdrawal period 10 days. Do not use in laying hens producing eggs for human consumption.

### PACKING:

1 Liter HDPE bottle.

