

# KYFOS E



Veterinary use



## Antibiotic combination - Powder for oral use

### FORMULA:

Each g of product contains:

Calcium Fosfomicin 300IU .....250 mg

Erythromycin .....100 mg

Excipients q.s.to..... 1 g

### DESCRIPTION:

Kyfos E is a powder to be administered in drinking water or mixed with feed. This product is a wide-spectrum antibiotic combination, prescribed for the treatment of avian infectious diseases caused by Mycoplasmas, Gram positive and Gram negative bacteria.

### PRESCRIPTION:

This product is indicated for the oral treatment of poultry, such as broiler chickens, breeders and replacement chickens. It is indicated in cases of mycoplasmosis, colibacillosis, chronic respiratory disease, staphylococcus-related infections, listeriosis, avian cholera, infectious coryza and diseases caused by Fosfomicin and/or Erythromycin-sensitive microorganisms.

### DOSAGE AND USE:

For oral administration in drinking water or feed.

**Poultry:** (30-40 mg/kg b.w.) In drinking water: 0.5 g per liter of drinking water. In Feed: Add 1 kg of product per ton of feed. Treat during 3 to 5 days.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

### WARNINGS:

Keep out of reach of children and domestic animals. Avoid inhalation and direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Store in original package in a cool and dry place between 15° and 25° C, protected against sunlight and moisture. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Validity after being mixed with feed: 14 days. Do not use after expiry date which is stated on the label.

Contraindications: none.

Side effects: very occasionally gastrointestinal disorders and other hypersensitivity reactions may occur. If you notice any serious side effects or other effects not listed on this notice, please inform your veterinarian.

Kyfos E has no contraindications or side effects at recommended doses.

**Use restrictions:** Withdrawal period in Poultry: 7 days. Do not use in laying hens producing eggs for human consumption.

### PACKING:

100 g, 1 kg and 5 kg sachets.

*May be possible that not all packages are traded.*

