

KYFOS T

Antibiotic combination - Powder for oral use



Veterinary use



FORMULA:

Each g of product contains:

Calcium Fosfomicin 300IU250 mg

Tylosin tartrate.....100 mg

Excipients q.s.to..... 1 g

DESCRIPTION:

Kyfos T is a wide-spectrum antibiotic combination, prescribed for the treatment of pigs and avian infectious diseases caused by Mycoplasmas, Gram positive and Gram negative bacteria. Among the most sensitive bacteria to the action of Fosfomicin, the enteropathogenic strains of Escherichia coli, the cause of colibacillosis, and Salmonella, typhoid and pullorosis agent are highlighted. Also its action on Pasteurella, Staphylococci, Streptococci and Pseudomonas is notable, and together with the addition of tylosin which is indicated to help, prevent, reduce injuries and lessen the severity of chronic respiratory disease of birds.

PRESCRIPTION:

This product is indicated for the oral treatment of poultry, such as broiler chickens, breeders and replacement chickens and pigs. It is indicated in cases of mycoplasmosis, colibacillosis, salmonellosis, chronic respiratory disease, staphylococcus-related infections, listeriosis, avian cholera, infectious coryza and diseases caused by Fosfomicin and/or Tylosin-sensitive microorganisms.

It is also indicated as treatment to prevent vibrio enteritis, dysentery and respiratory diseases in pigs.

DOSAGE AND USE:

For oral administration in drinking water or feed.

The dose for poultry and pigs is 100-110mg KYFOS T per kilogram of body weight.

Poultry:

In the feed: KYFOS T 1 kg / ton of feed.

In drinking water: approx 1 kg of KYFOS T / 1000 liters of water. Treat during 5 days.

Swines:

In feed: 2.5 kg KYFOS T / ton of feed.

In drinking water: 1.5 kg KYFOS T / 1000 liters of water.

Treat during 5 days.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS:

Keep out of reach of children and domestic animals. Avoid inhalation and direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Store in original package in a cool and dry place between 15° and 25° C, protected against sunlight and moisture. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Validity after being mixed with feed: 14 days. Do not use after expiry date which is stated on the label.

Contraindications: none.

Side effects: very occasionally gastrointestinal disorders and other hypersensitivity reactions may occur. May cause rectal mucosal edema, diarrhea and itching in piglets. If they happen to stop treatment immediately. If you notice any serious side effects or other effects not listed on this notice, please inform your veterinarian.

Kyfos T has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawal period in Poultry: 7 days. Pigs: 14 days. Do not use in laying hens producing eggs for human consumption.

PACKING:

100 g, 1 kg and 5 kg sachets.

May be possible that not all packages are traded.

