

KYXOFLOR 200P



Antibiotic - Hydrosoluble powder for oral use in drinking water

Veterinary use



FORMULA:

1 g of product contains:

Florfenicol200 mg

Excipients q. to..... 1 g

DESCRIPTION:

Kyxoflor 200P is a wide spectrum antibiotic with excellent absorption at intestinal level, broad safety margin, great resistance to bacterial inactivation and quick elimination from tissues, in the form of hydrosoluble powder.

PRESCRIPTION:

It is indicated for the curative treatment of enteric and respiratory infections caused by E coli and other florfenicol-sensitive bacteria, such as: coli enteritis, respiratory colibacillosis, colisepticaemia, E.coli-related complications of infectious diseases (coryza, CRD.).

For Porcines: Treatment and control of the Pneumonic complex caused by Actinobacillus pleuropneumoniae, Pasteurella spp and Haemophilus spp.

Gram negative: Pasteurella haemolytica, Pasteurella multocida. Haemophilus sommus, Haemophilus influenzae. Escherichia coli, Salmonella spp., Moraxella spp., Klebsiella spp., Enterococcus spp., Shigella spp., Treponema, Hyodysenteriae, Bacteroides spp., Bordetella bronchiseptica, Proteus spp. Neisseria spp.

Gram Positive: Corynebacterium pyogenes, Streptococcus agalactiae, Streptococcus dysgalactiae, Streptococcus uberis, Streptococcus zooepidermicus, Streptococcus aureus.

Target species: for pigs and poultry (broilers, replacement and breeder chickens).

DOSAGE AND USE:

Oral administration, dissolved in drinking water.

Poultry: 0.5 g of Kyxoflor 200P per liter of water (20 mg / kg body weight), for 3 to 5 days.

Pigs: 50 to 100 g of Kyxoflor 200P per 100 liters of water (10 - 20 mg / kg body weight), for 5 days.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS:

Keep out of the reach of children and domestic animals. Avoid inhalation and direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Store in original package in a cool and dry place between 15° and 25° C, protected against sunlight and moisture. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Keep adequate ventilation when administering product. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Do not use after expiry date which is stated on the label.

Contraindications: concurrent administration together with penicillins. Do not administer in pregnant sows or boars that are used for reproductive purposes.

Side effects: Diarrhea and / or erythema, edema, transient perianal and rectal can be observed, these effects can be observed during a week. If you notice any serious side effects or other effects not mentioned in this notice, inform your veterinarian.

Kyxoflor 200 P has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawal period: poultry 5 days and pigs 16 days. Do not use in laying hens producing eggs for human consumption.

PACKING:

500 g and 1 kg bags.

May be possible that not all packages are traded.

