

KYXOFLOR PREMIX



Veterinary use



Antibiotic - Premix for oral use in feed

FORMULA:

1 g of product contains:

Florfenicol300 mg

Excipients q.s.to..... 1 g

DESCRIPTION:

Kyxoflor Premix is a wide spectrum antibiotic with excellent absorption at intestinal level, broad safety margin, great resistance to bacterial inactivation and quick elimination from tissues.

PRESCRIPTION:

Curative treatment for enteric and respiratory infections caused by E coli and other florfenicol-sensitive bacteria, such as: coli enteritis, respiratory colibacillosis, colisepticaemia, E.coli-related complications of infectious diseases (coryza, CRD.).

For Porcines: Treatment and control of the Pneumonic complex caused by Actinobacillus pleuropneumoniae, Pasteurella spp and Haemophilus spp.

Gram negative: Pasteurella haemolytica, Pasteurella multocida. Haemophilus sommus, Haemophilus influenzae. Escherichia coli, Salmonella spp., Moraxella spp., Klebsiella spp., Enterococcus spp., Shigella spp., Treponema, Hyodysenteriae, Bacteroides spp., Bordetella bronchiseptica, Proteus spp. Neisseria spp.

Gram Positive: Corynebacterium pyogenes, Streptococcus agalactiae, Streptococcus dysgalactiae, Streptococcus uberis, Streptococcus zooepidermicus, Streptococcus aureus. Species: for porcines, broiler, replacement and breeder chickens.

DOSAGE AND USE:

Orally, mixed with feed.

Poultry (broilers, replacement chicks and breeders): 10-20 mg / kg of live weight (equivalent to 33.3 mg - 66.7 mg of Kyxoflor Premix per kg of weight), for 3 to 5 days.

Pigs: 2-10 mg / kg of live weight (equivalent to 6.7 mg - 33.3 mg of Kyxoflor Premix per kg of weight), for 5 consecutive days.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS:

Keep out of the reach of children and domestic animals. Avoid inhalation and direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Store in original package in a cool and dry place between 15° and 25° C, protected against sunlight and moisture. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Keep adequate ventilation when administering product. Close recipient tightly after every use. Validity after being mixed with feed: 15 days. Do not use after expiry date which is stated on the label.

Contraindications: concurrent administration together with penicillins. Do not administer in pregnant sows or boars that are used for reproductive purposes.

Side effects: Diarrhea and / or erythema, edema, transient perianal and rectal can be observed, these effects can be observed during a week. If you notice any serious side effects or other effects not mentioned in this notice, inform your veterinarian.

Kyxoflor Premix has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawal period: poultry 5 days and pigs 12 days. Do not use in laying hens producing eggs for human consumption.

PACKING:

5 kg and 20 kg bags.

May be possible that not all packages are traded.