

KYXOMOX 500



Veterinary use



Antibiotic - Powder for oral use

FORMULA:

Each g of product contains:

Amoxicillin*..... 500 mg

Excipients q.s.to..... 1 g

*(equivalent to 574 mg of Amoxicillin Trihydrate)

DESCRIPTION:

Soluble powder to be administered in drinking water or feed.

PRESCRIPTION:

This product contains of a wide spectrum antibiotic (Amoxicillin) with high oral absorption and broad tissue distribution . It is indicated to attack amoxicillin-sensitive infectious agents causing affections caused in the respiratory, genital-urinary and digestive systems. Its also prescribed for the treatment of bacterial complications of viral affections in poultry and pigs. Its antibacterial spectrum covers both gram-positive and gram-negative bacteria.

DOSAGE AND USE:

For oral administration in drinking water or feed at a rate of 20 mg of active material per kilogram of live weight for 3 to 5 days.

Multiply the amount of body weight to be treated by 0.04 to obtain the amount (in grams) of Kyxomox 500 to be used.

For poultry: Add 0,266 g of Kyxomox 500 / liters of water / day or 400 g of Kyxomox 500 / ton of feed / day.

For swine: Add 0.70 g of Kyxomox 500 / liters of water / day or 1.12 kg of Kyxomox 500 / ton of feed.

In the case of medicating the drinking water it is advisable to renew the water every 8 hours. Do not save for your future.

Instructions: the dose to be administered must be diluted before including it in the total volume of drinking water or feed: in 10 liters of water or in 10 kg of feed.

Consult the veterinarian to adjust the dose according to the needs of the case to be treated and according to the daily consumption of the flock.

WARNINGS:

Keep out of reach of children and pets. Avoid inhalation and direct contact of the product with the skin and eyes. If accidental exposure occurs, wash with running water and soap. Store in the original package in a cool and dry place between 15 ° and 25 ° C, protected from sunlight and moisture. People with known hypersensitivity to the drug should avoid contact with the product. Do not throw empty containers in open garbage dumps, dispose of them in accordance with local regulations. Close the container tightly after each use. It is advisable to prepare only the daily amount that will be consumed. Do not use solutions that have more than 12 hours of preparation.

The product remains stable once mixed with the food, for a period of 7 days. Do not use after the expiration date on the label.

Contraindications: hypersensitivity. Do not administer orally to rabbits, guinea pigs, equidae and hamsters since amoxicillin, like all aminopenicillins, has an important action on the caecal bacterial population.

Side effects: rashes, skin rashes and possible diarrhea and other hypersensitivity reactions. If you notice any serious side effects or other effects not mentioned in this notice, inform your veterinarian.

KYXOMOX 500 has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawal period in Poultry: 24 hours. Withdrawal period in pigs: 15 days.

Do not use laying hens producing eggs for human consumption.

PACKING:

400 g, 1 kg and 5 kg sachets.

May be possible that not all packages are traded.

