

KYXOTIL P



Veterinary use



Antibiotic - Powder for oral use

FORMULA:

1 g of product contains:

Tilmicosin (phosphate) 200 mg*

Excipients q.s. to..... 1 g

*(equivalent to 266 mg of tilmicosin phosphate)

DESCRIPTION:

KYXOTIL P is a medicinal premix for poultry and pigs containing Tilmicosin Phosphate to be administered in feed and drinking water.

PRESCRIPTION:

Tilmicosin is a macrolide antibiotic, synthesized from tylosin, to be used in pigs as feed additive for the control of respiratory diseases associated with: *Actinobacillus pleuropneumoniae*, *Pasteurella multocida* and *Mycoplasma hyopneumoniae*. Tilmicosin has proven to be successful against *Mycoplasma gallisepticum* in poultry. Can also be administered in drinking water. Tilmicosin is intercepted by macrophage and polymorphonuclear and transported to the infected zone, reaching high concentrations for long periods of time. Its pharmacokinetic-pharmacodynamic action (antibacterial action, dependent concentration, good liposolubility, high tissue distribution, penetration and intracellular distribution) is particularly beneficial as it can be administered as single dose thus making it into a therapeutic tool against pulmonary, ocular and foot bacterial infections. Its wide therapeutic spectrum acts significantly against both Gram-positive and certain Gram-negative bacteria and mycoplasma: *Pasteurella haemolytica*, *Pasteurella multocida*, *Haemophilus somnus*, *Mycoplasma dispar*, *Mycoplasma bovirhinis*, *Staphylococcus aureus*, *Streptococcus agalactiae*, *Actinomyces pyogenes*, *Moraxella boris*, *Actinobacillus pleuropneumoniae*, *Erysipelothrix rhusiopathiae*, *Clostridium perfringens*, *Fusobacterium necrophorum*.

DOSAGE AND USE:

Oral administration.

Porcines: The dose is 10 to 20 mg /kg of b.w. per day, equivalent to approx 1 g to 2 g of product per kg of feed. Equivalent of same dose for use in drinking water is approximately 0,87 g to 1,7 g per liter of drinking water. For specific cases:

Dysentery treatment: In Feed: 500 g / ton. In Drinking water: 60 g / 100 L. Duration: 3 to 7 days.

Control and prevention of dysentery: In Feed: 200 g / ton. In Drinking water: 25 g / 100 L. Duration: During risk period.

Enzootic pneumonia: In Feed: 500 g / ton. In Drinking water: 60 g / 100 L. Duration: 7 days.

Other cases: Continuous treatment during 10 - 21 days.

Poultry: The dose is 10 to 15 mg/kg of b.w. per day, equivalent to 40 g to 55 g or up to 100 g of product/ 100 liters of drinking water.

Equivalent of same dose for use in feed is approx. 375 g to 550 g or up to 1 kg of product per ton of feed.

Continuous treatment during 5 to 6 days.

In order to ensure the ingestion of the recommended dose, the medicated water will be the only drinking water to be supplied.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS:

The product must be stored in its original container at a temperature between 15° and 25°C. Keep out of the reach of children. Avoid direct contact with skin and eyes. Persons showing known hypersensitivity to this drug must avoid contact with product. The operator must wear mask, eye goggles, rubber gloves and overall. Keep adequate ventilation when administering product. Handle with care on pregnant animals and those destined for rearing. Once mixed with feed, duration of product is 15 days. For drinking water use, it is recommended to prepare the daily quantity to be consumed. Do not use solutions whose time of preparation exceeds 24 hours.

Do not throw empty containers in open garbage dump sites, discard them according to local regulations.

Contraindications: Do not allow horses and other equines to drink water containing tilmicosin. Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

Side effects: In very rare cases, a decrease in water consumption has been observed. If you notice any serious side effects or other effects not listed on this notice, please inform your veterinarian.

KYXOTIL P has no contraindications or side effects at recommended doses.

Use restrictions: withdrawal period: 7 days of the last treatment. Do not use in laying hens producing eggs for human consumption.

PACKING:

1 kg, 5 kg aluminium sachets and 20 kg bags.

May be possible that not all packages are traded.

KYXON
PHARMACEUTICALS S.R.L.



Av. Pedro Goyena 1432, 1°A
Buenos Aires, Argentina.
Email: info@kyxon.com.ar