

KYXOTIL S



Antibiotic - Solution for oral use

FORMULA:

1 ml of product contains:
Tilmicosin (as phosphate)* 250 mg
Vehicle q.s.to 1 ml
**equivalen to 332 mg of tilmicosin phosphate.*

Veterinary use



DESCRIPTION:

KYXOTIL S is a solution containing Tilmicosin phosphate to be administered in drinking water of poultry, pigs and calves.

PRESCRIPTION:

Tilmicosin is a macrolide antibiotic, synthesized from tylosin. For use in the symptomatic and specific treatment of the respiratory diseases associated with *Pasteurella* and *Mycoplasma* species. Swine: For treatment of respiratory diseases associated to *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and other organisms sensible to Tilmicosin. Poultry: For treatment of respiratory diseases associated to *Mycoplasma gallisepticum* and *Mycoplasma synoviae*. Calves: For treatment of respiratory diseases in calves associated to *Mycoplasma haemolytica*, *Pasteurella multocida*, *Mycoplasma bovis*, *Mycoplasma dispar* and other organisms sensible to tilmicosin. Tilmicosin is a wide spectrum antibiotic effective against Gram positive, Gram negative bacteria and mycoplasmas: *Pasteurella haemolytica*, *Pasteurella multocida*, *Haemophilus somnus*, *Mycoplasma dispar*, *Mycoplasma bovirhinis*, *Staphylococcus aureus*, *Streptococcus agalactiae*, *Actinomyces pyogenes*, *Actinobacillus pleuropneumoniae*, *Erysipelothrix rhusiopathiae*, *Moraxella boris*, *Clostridium perfringens*, *Fusobacterium necrophorum*.

DOSAGE AND USE:

For oral administration in drinking water.

Poultry: The dose is 10 to 15 mg / kg per day, equivalent to 50 to 75 mg of Tilmicosin/ liter of drinking water, that is 20 to 30 ml of product/ 100 liter of drinking water, continuous treatment during 5 days.

Swine: The dose is 10 to 20 mg /kg per day, equivalent to 175 to 350 mg of Tilmicosin/liter of drinking water, that is 70 to 140 ml per 100 liter of drinking water, continuous treatment during 10 days.

Calves: The dose is 12.5 mg/kg b.w. equivalent to 1 ml of product/ 20 kg b.w., continuous treatment during 3 to 5 days. Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS: Keep out of the reach of children and domestic animals. Avoid direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Use suitable and protective garment when administering medication. Wash hands and change clothes after administration. Store in its original container at temperatures between 15°C and 25°C avoiding direct sunlight on stored product. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Do not freeze. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Do not use after expiry date which is stated on the label. Use with care in pregnant animals or those confined for rearing.

Contraindications: Do not allow horses and other equines to drink water containing tilmicosin. Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

Side effects: In very rare cases, a decrease in water consumption has been observed. If you notice any serious side effects or other effects not listed on this notice, please inform your veterinarian.

KYXOTIL S has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawals: Porcines: 14 days. Poultry: 12 days. Calves: 42 days.

Do not administer in laying hens producing eggs for human consumption.

PACKING:

240 mL and 1 Liter HDPE bottle.

May be possible that not all packages are traded.

