

KYXURIL

Internal Antiparasitic - Coccidicide - Solution for oral use



Veterinary use



FORMULA:

Each ml of product contains:

Diclazuril.....10 mg

Vehicle q.s.to..... 1 ml

DESCRIPTION:

KYXURIL is an oral solution based on diclazuril which acts as internal antiparasitic.

PRESCRIPTION:

KYXURIL is an anticoccidial intended for treatment and prevention of Coccidiosis caused by Eimerias in poultry, ovines, porcines and rabbits. KYXURIL completely interrupts the life cycle of coccidial microorganisms. It exerts a powerful action against several stages: oocysts, schizont, merozoite, macro and micro gametocytes. Interrupts the life cycle of microorganisms in early stages. KYXURIL acts on the following coccidials: Eimeria tenella (cecum of the chicken and hens), Eimeria mivati (small intestine of chickens), Eimeria necatrix (small intestine and hen cecum), Eimeria brunetti (small intestine of chickens), Eimeria maxima (small intestine of chickens), Eimeria acervulina (small intestine of chickens), Eimeria mitis (small intestine of chickens), Eimeria ovina (small intestine of sheep), Eimeria suis (small intestine of pigs), Eimeria stiedae (small intestine and rabbit cecum).

To treat porcines, poultry, rabbits and ovines.

DOSAGE AND USE:

For oral administration in drinking water.

Poultry: dose is 1 mg/kg/body weight. Assuming that water consumption is between 120 and 200 ml/kg, 500 to 830 ml of product should be diluted every 1000 liters of drinking water. This is an estimated dose and must be adjusted according to production and seasonal variations in water consumption on each farm. Administer continuously for 48 hours.

Ovines: 20 mg/kg/day, continuous treatment for 48 hours

Rabbits: 2 to 4 ppm, continuous treatment for 48 hours.

Porcines: 5 mg/kg/body weight, only one dose.

Consult Veterinary Doctor in order to adjust dose according to the needs of the case to be treated and according to daily flock consumption.

WARNINGS:

Keep out of the reach of children and domestic animals. Avoid direct contact of product with skin and eyes. If accidental exposure occurs, wash with running water and soap. Use suitable garment when administering medication. Wash hands and change clothes after administration. Store in its original container at temperatures between 15°C and 25°C avoiding direct sunlight on stored product. People with known hypersensitivity to the drug must avoid contact with product. Do not throw empty containers in open garbage dump sites, discard them according to local regulations. Do not freeze. Close recipient tightly after every use. It is advised to prepare only the daily quantity to be consumed. Do not use solutions exceeding 24 hours from preparation. Do not use after expiry date which is stated on the label. Use with care in pregnant animals or those confined for rearing.

Contraindications: in case of hypersensitivity.

Side effects: If you notice any serious side effects, please inform your veterinarian.

KYXURIL has no contraindications or side effects at recommended doses.

Use restrictions: Withdrawal period is 5 days for all species. Do not administer in laying hens producing eggs for human consumption.

PACKING:

250 ml and 1 Liter HDPE bottle.

May be possible that not all packages are traded.

